

STANDARD TREATMENT WORKFLOW (STW)

ACUTE ENCEPHALITIS SYNDROME –(AES) IN CHILDREN

Shally Awasthi¹, Sushil Kabra², Neelam Mohan³, Pushpa Kini⁴, Suvasini Sharma⁵, Joseph Mathew⁶, Surjit Singh⁷, Kuldeep Singh⁸, Himanshu Chaturvedi⁹, Shinjini Bhatnagar¹⁰

¹King George's Medical University, Lucknow; ²All India Institute Of Medical Science, New Delhi; ³Medanta, Gurgaon; ⁴Kasturba Medical College, Manipal; ⁵Lady Hardinge Medical College, New Delhi; ⁶Postgraduate Institute of Medical Education and Research, Chandigarh; ⁷Postgraduate Institute of Medical Education and Research, Chandigarh; ⁸All India Institute Of Medical Science, Jodhpur; ⁹Balrampur Hospital, Lucknow; ¹⁰Translational Health Science and Technology Institute (THSTI), Faridabad

CORRESPONDING AUTHOR

Dr. Shally Awasthi, Professor & Head, Department of Paediatrics, King George's Medical University, Lucknow Uttar Pradesh 226003

Email: shally07@gmail.com

CITATION

Awasthi S, Kabra S, Mohan N, Kini P, Sharma S, Mathew J, Singh S, Singh K, Chaturvedi H, Bhatnagar S. ACUTE ENCEPHALITIS SYNDROME –(AES) IN CHILDREN. Journal of the Epidemiology Foundation of India. 2024;2(1Suppl):S01-S02. DOI: <https://doi.org/10.56450/JEFI.2024.v2i1Suppl.001>

This work is licensed under a Creative Commons Attribution 4.0 International License.

©The Author(s). 2024 Open Access

DISCLAIMER

This article/STW, was originally published by Indian Council of Medical Research (ICMR) under Standard Treatment Workflow. The reprinting of this article in Journal of the Epidemiology Foundation of India (JEFI) is done with the permission of ICMR. The content of this article is presented as it was published, with no modifications or alterations. The views and opinions expressed in the article are those of the authors and do not necessarily reflect the official policy or position of JEFI or its editorial board. This initiative of JEFI to reprint STW is to disseminate these workflows among Health Care Professionals for wider adoption and guiding path for Patient Care.



Standard Treatment Workflow (STW) for the Management of ACUTE ENCEPHALITIS SYNDROME (AES) IN CHILDREN ICD-10-G04

October 2019

Acute onset of fever ($\leq 5-7$ days) with altered sensorium and/or new onset of seizures (excluding simple febrile seizures)

SYMPTOMS

- Fever, headache, vomiting, lethargy, unconsciousness
- Seizures
- Abnormal posturing
- Paucity of limb movements

ADDITIONAL INFORMATION (HISTORY OF)

- Rash, vesicles, past history of chicken pox
- Residence of child: rural/urban, endemic for cerebral malaria, any epidemic of AES in neighborhood
- Animal contact, insect bite, dog bite
- Drug or toxin exposure: enquire for presence of any drugs at home
- Recent travel
- Trauma
- Seizures
- Recent immunizations
- Recurrent episodes of encephalopathy
- Past or concurrent systemic illness
- Pre-morbid developmental/ neurological status of the child

EXAMINATION		
VITAL SIGNS	GENERAL EXAMINATION	NEUROLOGICAL EXAMINATION
<ul style="list-style-type: none"> Temperature Pulse rate Respiratory rate Blood pressure 	<ul style="list-style-type: none"> Pallor Petechiae Rash Icterus 	<ul style="list-style-type: none"> Level of consciousness by Glasgow Coma Scale (GCS) Abnormal posturing: decerebrate, decorticate Active seizures Cranial nerves, pupil size and reaction, doll's eye movements, squint, facial deviation Focal neurological deficits Meningeal signs

INVESTIGATIONS		
ESSENTIAL	DESIRABLE	OPTIONAL
CBC, LFT, KFT, blood sugar, CECT Brain, CSF examination* (cytology, biochemistry, culture, AFB staining, Gene Xpert), peripheral smear for malarial parasite, Rapid Malarial Antigen Test	MRI Brain, CSF PCR for Herpes simplex encephalitis, JE serology, EEG, Dengue serology and NSI testing, HIV testing	CSF Neurovirology panel, anti-NMDA receptor antibody testing, PCR viral testing of other samples (throat swab, nasopharyngeal aspirates, stool etc), Blood Tandem Mass Spectrometry and urine gas chromatography, antinuclear antibodies

*Lumbar puncture is contra-indicated or neuroimaging must be obtained before lumbar puncture
 1.Fundus: papilledema 2. Platelet count < 50,000 3. Focal neurological deficits 4. Asymmetric/unreactive pupils 5.Decerebrate/decorticate posturing

MANAGEMENT

All patients need to be admitted.
 If any of the following signs are present, the child should be referred to tertiary care facility with PICU and facilities for mechanical ventilation:
 • Glasgow Coma Scale < 8 • Abnormal breathing pattern • Shock not responding to fluid bolus • Decerebrate or decorticate posturing
 • Seizures persisting despite benzodiazepine and phenytoin

Step I: Rapid assessment and stabilization

- Establish and maintain airway: Intubate if GCS<8, impaired airway reflexes, abnormal respiratory pattern, signs of raised intracranial pressure, SpO₂ <92% despite high flow oxygen and fluid refractory shock
- Ventilation, oxygenation
- Circulation: Establish IV access, take samples for relevant investigations, fluid bolus if in circulatory failure (20 mL/kg NS), inotropes if required
- Identify signs of cerebral herniation or raised ICP
- Temperature: treat fever and hypothermia
- Treat ongoing seizures- Benzodiazepine, followed by phenytoin loading

Step II: History, Examination and Investigations as given above

Step III: Empirical Treatment (must be started if CSF cannot be done/ report will take time and patient sick)

- Ceftriaxone: 100 mg/kg/day in 2 divided doses X 10-14 days
- Acyclovir (use in all suspected sporadic viral encephalitis): 3 mo to 12 y: 500mg/m² 8 hourly (min 21 days) >12 y: 10mg/kg 8 hourly (14-21 days in confirmed cases)**
- Artesunate combination therapy (stop if peripheral smear and RDT are negative): 3mg/kg in child <20 kg, and 2.4mg/kg in child >20kg IV/IM at 0,12 and 24 hours, followed by once daily parental/oral X 3-7 days

**If therapy was started empirically stop acyclovir, in case an alternative diagnosis is confirmed, or HSV PCR of CSF is negative on two occasions (24-48 h apart) and MRI imaging not suggestive of Herpes Simplex Encephalitis

Step IV: Supportive care and treatment

- Maintain euglycemia, hydration and control fever
- Treat raised intracranial pressure#, mild head-end elevation-15-30°
- Treat seizures###: Give anticonvulsant if: history of seizures / GCS <8 / child has features of raised ICP
- Steroids: Pulse steroids (methylprednisolone) to be given in children with suspected acute disseminated encephalomyelitis or autoimmune encephalitis

Step V: Prevention/treatment of complications and rehabilitation

- Physiotherapy, posture change, prevent bed sores and exposure keratitis
- Complications: aspiration pneumonia, nosocomial infections, coagulation disturbances
- Nutrition: early feeding
- Psychological support to patient and family

<h4>#Management of raised intracranial pressure</h4> <ul style="list-style-type: none"> Intubate if: GCS <8 / evidence of herniation / irregular respirations and inability to maintain airway Signs of impending herniation: patient to be hyperventilated to a target PaCO₂ of 30-35 mmHg Initial bolus of Mannitol(0.25 g/kg), then 0.25 g/kg q 6 h as per requirement, up to 48 hours In the presence of hypotension, hypovolemia, and renal failure: hypertonic (3%) saline (preferable to mannitol) 0.1-1 mL/kg/hr by infusion; serum sodium to be targeted to 145-155 meq/L Adequate sedation and analgesia Avoid noxious stimuli Administer nebulized lignocaine prior to endotracheal tube suctioning 	<h4>##Treatment of seizures</h4> <p>1st Line: IV Lorazepam 0.1mg/kg or Midazolam 0.2 mg/kg or Diazepam 0.3 mg/kg).</p> <p>If no IV access: IM Midazolam 0.2 mg/kg</p> <p>2nd Line: Inj. Phenytoin 20 mg/kg (in Normal saline 1mg/kg/min)</p> <p>If seizures still persist: Refractory status: Transfer to PICU -> midazolam infusion (1-18 microgram/kg/min) If ICU facilities not available, sodium valproate (20 mg/kg) or levetiracetam (20-40 mg/kg) or phenobarbitone (20mg/kg)</p>
---	---

DISCHARGE CRITERIA

Hemodynamically stable	Improvement in consciousness	Afebrile	Has started eating and drinking orally	Seizures have subsided	Parents have been explained the supportive care and physiotherapy to be continued at home
------------------------	------------------------------	----------	--	------------------------	---

➔ KEEP A HIGH THRESHOLD FOR INVASIVE PROCEDURES

REFERENCES

- World Health Organisation. Acute Encephalitis Syndrome. Japanese encephalitis surveillance standards. January 2006. From WHO-recommended standards for surveillance of selected vaccine-preventable diseases. WHO/VIS/03.01. Available from: <http://www.who.int/vaccines-documents/DocxPDF06/843.pdf>
- National Program for Prevention and Control of Japanese Encephalitis/Acute Encephalitis Syndrome 2014. Government of India Ministry of Health & Family Welfare Directorate General of Health Services National Vector Borne Disease Control Programme.
- Sharma S, Mishra D, Aneja S, Kumar R, Jain A, Vashishtha VM. Consensus guidelines on evaluation and management of suspected acute viral encephalitis in children in India. Indian Pediatr. Nov 2012;49(11):897-910.
- Sankhyani N, Vykunta Raju KN, Sharma S, Gulati S. Management of raised intracranial pressure. Indian J Pediatr. 2010 Dec;77(12):1409-16.

This STW has been prepared by national experts of India with feasibility considerations for various levels of healthcare system in the country. These broad guidelines are advisory, and are based on expert opinions and available scientific evidence. There may be variations in the management of an individual patient based on his/her specific condition, as decided by the treating physician. There will be no indemnity for direct or indirect consequences. Kindly visit our web portal (stw.icmr.org.in) for more information.
 © Indian Council of Medical Research and Department of Health Research, Ministry of Health & Family Welfare, Government of India.