

STANDARD TREATMENT WORKFLOW (STW)

Hysterectomy for Benign Gynaecological Conditions

Reva Tripathi¹, Vinita Das², Manju Puri³, Radhika⁴, Neelam Aggarwal⁵, Asmita Rathore⁶, Aruna Kekre⁷, Dasari Papa⁸, Usha Rani⁹, Manika Khanna¹⁰, Neerja Bhatla¹¹, Seema Saran¹²
¹Maulana Azad Medical College (MAMC), New Delhi; ²King George's Medical College, Lucknow; ³Lady Hardinge Medical College, New Delhi; ⁴University College of Medical Sciences, New Delhi; ⁵Postgraduate Institute of Medical Education and Research, Chandigarh; ⁶Maulana Azad Medical College (MAMC), New Delhi; ⁷Christian Medical College, Vellore; ⁸Jawaharlal Institute of Postgraduate Medical Education & Research, Puducherry; ⁹Institute of Obstetrics and Gynaecology, Chennai; ¹⁰NRIGS; ¹¹All India Institute Of Medical Science, New Delhi; ¹²Government Medical College Budaun

CORRESPONDING AUTHOR

Dr. Reva Tripathi, Department of OBS/GYN, Maulana Azad Medical College (MAMC), New Delhi.

Email: revatripathi@gmail.com

CITATION

Tripathi R, Das V, Puri M, Radhika, Aggarwal N, Rathore A, Kekre A, Papa D, Rani U, Khanna M, Bhatla N, Saran S. Hysterectomy for Benign Gynaecological Conditions. Journal of the Epidemiology Foundation of India. 2024;2(1Suppl):S41-S42. DOI: <https://doi.org/10.56450/JEFI.2024.v2i1Suppl.021>

This work is licensed under a Creative Commons Attribution 4.0 International License.

@The Author(s). 2024 Open Access

DISCLAIMER

This article/STW, was originally published by Indian Council of Medical Research (ICMR) under Standard Treatment Workflow. The reprinting of this article in Journal of the Epidemiology Foundation of India (JEFI) is done with the permission of ICMR. The content of this article is presented as it was published, with no modifications or alterations. The views and opinions expressed in the article are those of the authors and do not necessarily reflect the official policy or position of JEFI or its editorial board. This initiative of JEFI to reprint STW is to disseminate these workflows among Health Care Professionals for wider adoption and guiding path for Patient Care.


October/2019



Standard Treatment Workflow (STW) for HYSTERECTOMY FOR BENIGN GYNAECOLOGICAL CONDITIONS

IN WOMEN AGED LESS THAN 40 AND/OR LOW PARITY IT IS **MANDATORY** TO HAVE A SECOND OPINION FROM A QUALIFIED GYNAECOLOGIST


HYSTERECTOMY TO BE CONSIDERED ONLY WHEN CHILD BEARING IS COMPLETED & RARELY IN YOUNGER PATIENTS



Standard Treatment Workflow (STW) for HYSTERECTOMY FOR BENIGN GYNAECOLOGICAL CONDITIONS

IN WOMEN AGED LESS THAN 40 AND/OR LOW PARITY IT IS **MANDATORY** TO HAVE A SECOND OPINION FROM A QUALIFIED GYNAECOLOGIST

HYSTERECTOMY TO BE CONSIDERED ONLY WHEN CHILD BEARING IS COMPLETED & RARELY IN YOUNGER PATIENTS



HYSTERECTOMY SHOULD NOT BE DONE FOR

White discharge per vaginum	Cervicitis	Non specific abdominal or pelvic pain	Minor degree of utero vaginal prolapse	Fibroids which are small (less than 5 cm) or Asymptomatic (less than 12 weeks size uterus)	Simple ovarian cyst less than or equal to 5 cm
-----------------------------	------------	---------------------------------------	--	--	--

COMPONENTS OF PRE OPERATIVE COUNSELLING AND INFORMED CONSENT

- Need for hysterectomy
- Alternative treatment options
- Risks and benefits
- Potential complications of the procedure
- Removal/ conservation of ovaries & tubes
- Route of hysterectomy
- Possible need for post operative Hormone therapy in selected cases

INVESTIGATIONS

- Complete Blood Count
- Blood grouping & cross matching
- Fasting Blood Sugar & Post Prandial Blood Sugar
- Renal Function Test
- Liver Function Test
- Urine Routine & Microscopy
- Electrocardiogram
- X ray chest
- Others as indicated

COMPLICATIONS TO BE EXPLAINED

- Risk of Infection
- Bleeding (primary/ reactionary/ secondary)
- Injury to bladder/ bowel/ ureter
- Pain
- Fever
- Hernia (rare and late complication)

FOLLOW UP

- Discharge summary with operative details
- Review for histopathology report
- Report if there is fever, bleeding or any other symptoms
- Avoid lifting heavy weight for 8 weeks
- Abstinence for eight weeks
- Adequate iron and calcium & Vitamin D3 supplements
- Evaluate need for hormones in very selected patients

Ovaries should be preserved in most pre-menopausal women unless diseased or removal specifically indicated

While doing hysterectomy for benign gynaecological conditions in pre-menopausal women, it is recommended to combine it with bilateral salpingectomy with a view to minimise the risk of subsequent development of ovarian malignancy^{1,2}

1. Pérez-López FR et al. Interventions to reduce the risk of ovarian and fallopian tube cancer: A European Menopause and Andropause Society Position Statement. *Maturitas*. 2017
2. Dorellus A et al. Efficacy of salpingectomy at hysterectomy to reduce the risk of epithelial ovarian cancer: a systematic review. *BJOG*. 2017.

COUNSELLING IS AN IMPORTANT ADJUNCT TO MANAGEMENT

KEEP A HIGH THRESHOLD FOR INVASIVE PROCEDURES

This STW has been prepared by national experts of India with feasibility considerations for various levels of healthcare system in the country. These broad guidelines are advisory, and are based on expert opinions and available scientific evidence. There may be variations in the management of an individual patient based on his/her specific condition, as decided by the treating physician. There will be no indemnity for direct or indirect consequences. Kindly visit our web portal (stw.icmr.org.in) for more information.
© Indian Council of Medical Research and Department of Health Research, Ministry of Health & Family Welfare, Government of India.