

STANDARD TREATMENT WORKFLOW (STW)

Uterine Fibroids and Polyps

Reva Tripathi¹, Vinita Das², Manju Puri³, Radhika⁴, Neelam Aggarwal⁵, Asmita Rathore⁶, Aruna Kekre⁷, Dasari Papa⁸, Usha Rani⁹, Manika Khanna¹⁰, Neerja Bhatla¹¹, Seema Saran¹²

¹Maulana Azad Medical College (MAMC), New Delhi; ²King George's Medical College, Lucknow; ³Lady Hardinge Medical College, New Delhi; ⁴University College of Medical Sciences, New Delhi; ⁵Postgraduate Institute of Medical Education and Research, Chandigarh; ⁶Maulana Azad Medical College (MAMC), New Delhi; ⁷Christian Medical College, Vellore; ⁸Jawaharlal Institute of Postgraduate Medical Education & Research, Puducherry; ⁹Institute of Obstetrics and Gynaecology, Chennai; ¹⁰NRIGS; ¹¹All India Institute Of Medical Science, New Delhi; ¹²Government Medical College Budaun

CORRESPONDING AUTHOR

Dr. Reva Tripathi, Department of OBS/GYN, Maulana Azad Medical College (MAMC), New Delhi.

Email: revatripathi@gmail.com

CITATION

Tripathi R, Das V, Puri M, Radhika, Aggarwal N, Rathore A, Kekre A, Papa D, Rani U, Khanna M, Bhatla N, Saran S. Uterine Fibroids and Polyps. Journal of the Epidemiology Foundation of India.

2024;2(1Suppl):S45-S46. DOI: <https://doi.org/10.56450/JEFI.2024.v2i1Suppl.023>

This work is licensed under a Creative Commons Attribution 4.0 International License.

©The Author(s). 2024 Open Access


DISCLAIMER

This article/STW, was originally published by Indian Council of Medical Research (ICMR) under Standard Treatment Workflow. The reprinting of this article in Journal of the Epidemiology Foundation of India (JEFI) is done with the permission of ICMR. The content of this article is presented as it was published, with no modifications or alterations. The views and opinions expressed in the article are those of the authors and do not necessarily reflect the official policy or position of JEFI or its editorial board. This initiative of JEFI to reprint STW is to disseminate these workflows among Health Care Professionals for wider adoption and guiding path for Patient Care.


0102 / JEFI 2024




Standard Treatment Workflow (STW) for the Management of UTERINE FIBROIDS AND POLYPS ICD-10-D25 & N84



**Standard Treatment Workflow (STW) for the Management of
UTERINE FIBROIDS AND POLYPS**
ICD-10-D25 & N84



ASYMPTOMATIC FIBROIDS <5CM DO NOT NEED TO BE TREATED



*Norethisterone (max daily dose 40 mg) OR Medroxyprogesterone acetate (max daily dose 40 mg). Any hormone should be given orally daily in divided doses for a duration of three weeks and repeated in a cyclical manner for total of 4-6 cycles of treatment

ALL THERAPUTIC OPTIONS NEED TO BE EXPLAINED TO THE PATIENT INCLUDING JUST KEEPING THE PATIENT ON OBSERVATION. ALL PATIENTS OF FIBROID UTERUS DO NOT NECESSARILY NEED HYSTERECTOMY.

👉 COUNSELLING IS AN IMPORTANT ADJUNCT TO MANAGEMENT

KEEP A HIGH THRESHOLD FOR INVASIVE PROCEDURES

This STW has been prepared by national experts of India with feasibility considerations for various levels of healthcare system in the country. These broad guidelines are advisory, and are based on expert opinions and available scientific evidence. There may be variations in the management of an individual patient based on his/her specific condition, as decided by the treating physician. There will be no indemnity for direct or indirect consequences. Kindly visit our web portal (stw.icmr.org.in) for more information. © Indian Council of Medical Research and Department of Health Research, Ministry of Health & Family Welfare, Government of India.