

STANDARD TREATMENT WORKFLOW (STW)

Adult Pericardial Tuberculosis

Dhruva Chaudhry¹, Ashutosh N Aggarwal², Anil Kumar Jain³, Ashwani Khanna⁴, Camilla Rodrigues⁵, Jai Bhagwan Sharma⁶, Jyotirmay Biswas⁷, Kusum Sharma⁸, Mandira Varma-Basil⁹, Manish Modi¹⁰, Manjula Datta¹¹, Narayan Jana¹², Nitish Naik¹³, Priscilla Rupali¹⁴, Rajesh Malhotra¹⁵, Ramprasad Dey¹⁶, Ritesh Aggarwal¹⁷, Rohit Bhatia¹⁸, Roy Thankachen¹⁹, Sambit N Bhattacharya²⁰, Thangakunam Balamugesh²¹, Uday Pratap Singh²², V Ramesh²³, Vineet Ahuja²⁴, Vishal Sharma²⁵, Vishali Gupta²⁶

¹Pulmonary & Critical Care Medicine, Pandit Bhagwat Dayal Sharma Post Graduate Institute of Medical Sciences, Rohtak; ²Pulmonary Medicine, Postgraduate Institute of Medical Education and Research, Chandigarh; ³Orthopedics, University College of Medical Sciences, New Delhi; ⁴National Tuberculosis Elimination Program, Govt of India, New Delhi; ⁵Parmanand Deepchand Hinduja and Medical Research Centre, Mumbai; ⁶Obstetrics and Gynaecology, All India Institute of Medical Sciences, New Delhi; ⁷Uveitis & Ocular Pathology Department, Sankara Nethralaya, Chennai; ⁸Medical Microbiology, Postgraduate Institute of Medical Education and Research, Chandigarh; ⁹Microbiology, Vallabhbhai Patel Chest Institute, University of Delhi, Delhi; ¹⁰Neurology, Postgraduate Institute of Medical Education and Research, Chandigarh; ¹¹ASPIRE Chennai; ¹²Obstetrics and Gynaecology, Chittaranjan Seva Sadan College of Obstetrics, Gynaecology and Child Health, Kolkata; ¹³Cardiology, All India Institute of Medical Sciences, New Delhi; ¹⁴Infectious Diseases, Christian Medical College, Vellore; ¹⁵Orthopedics, All India Institute of Medical Sciences, New Delhi; ¹⁶Obstetrics and Gynaecology, Chittaranjan Seva Sadan College of Obstetrics, Gynaecology and Child Health, Kolkata; ¹⁷Pulmonary Medicine, Postgraduate Institute of Medical Education and Research, Chandigarh; ¹⁸Neurology, All India Institute of Medical Sciences, New Delhi; ¹⁹Cardio-thoracic and Vascular Surgery, Christian Medical College Vellore; ²⁰Dr Baba Saheb Ambedkar Medical College & Hospital, Delhi; ²¹Pulmonary Medicine, Christian Medical College, Vellore; ²²Urology, Sanjay Gandhi Postgraduate Institute of Medical Sciences, Lucknow, India; ²³Employees' State Insurance Corporation Medical College and Hospital, Faridabad; ²⁴Gastroenterology, All India Institute of Medical Sciences, New Delhi; ²⁵Gastroenterology, Postgraduate Institute of Medical Education and Research, Chandigarh; ²⁶Advanced Eye Centre, Postgraduate Institute of Medical Education and Research, Chandigarh

CORRESPONDING AUTHOR

Dhruva Chaudhry, Pulmonary & Critical Care Medicine, Pandit Bhagwat Dayal Sharma Post Graduate Institute of Medical Sciences, Rohtak, Haryana

Email: dhruvachaudhry@yahoo.co.in

CITATION

Chaudhry D, Aggarwal AN, Jain AK, Ashwani Khanna, Rodrigues C, Sharma JB, Biswas J, Sharma K, Varma M, Modi M, Datta M, Jana N, Naik N, Rupali P, Malhotra R, Dey R, Aggarwal R, Bhatia R, Thankachen R, Bhattacharya SN, Balamugesh T, Singh UP, Ramesh V, Ahuja V, Sharma V, Gupta V. Adult Pericardial Tuberculosis. Journal of the Epidemiology Foundation of India. 2024; 2(1Suppl):S221-S222.

DOI: <https://doi.org/10.56450/JEFI.2024.v2i1Suppl.111>


This work is licensed under a Creative Commons Attribution 4.0 International License.

©The Author(s). 2024 Open Access


DISCLAIMER

This article/STW, was originally published by Indian Council of Medical Research (ICMR) under Standard Treatment Workflow. The reprinting of this article in Journal of the Epidemiology Foundation of India (JEFI) is done with the permission of ICMR. The content of this article is presented as it was published, with no modifications or alterations. The views and opinions expressed in the article are those of the authors and do not necessarily reflect the official policy or position of JEFI or its editorial board. This initiative of JEFI to reprint STW is to disseminate these workflows among Health Care Professionals for wider adoption and guiding path for Patient Care.

March 2022



Department of Health Research
Ministry of Health and Family Welfare, Government of India



INDIAN COUNCIL OF
MEDICAL RESEARCH
Serving the nation since 1951

Standard Treatment Workflow (STW) for the Management of ADULT PERICARDIAL TUBERCULOSIS ICD-10-A18.84

WHEN TO SUSPECT	COMPLICATIONS
<p>SYMPTOMS</p> <ul style="list-style-type: none"> • Cough, fever, breathlessness or pleuritic chest pain • May be associated with weight loss, night sweats or difficulty lying down • Past history or a history of contact with a patient with a diagnosis of tuberculosis • Examination reveals tachycardia, increased jugular venous pressure, hepatomegaly, ascites, & peripheral edema • A pericardial friction rub and distant heart sounds present on cardiovascular examination • If clinical picture +/- heart US suggest pericarditis or pericardial effusion refer for echo-cardiogram 	<p>Constrictive pericarditis: Clinical signs for recognition include</p> <ul style="list-style-type: none"> • Kussmaul's sign (lack of an inspiratory decline in jugular venous pressure) • Elevated & distended jugular veins with a prominent Y descent (second inward deflection of internal jugular pulse due to diastolic inflow of blood into the right ventricle) • Pericardial knock (rare) <p>Cardiac tamponade: Clinical signs include</p> <ul style="list-style-type: none"> • Sinus tachycardia • Hypotension with a narrow pulse pressure • Elevated JVP jugular venous pressure • Muffled heart sounds • Pulsus paradoxus (a decrease in systolic blood pressure by >10 mmHg on inspiration) • Ascites <p>Other complications:</p> <ul style="list-style-type: none"> • Myopericarditis: Abnormal ejection fraction with evidence of myocarditis and pericarditis (elevated cardiac enzymes & ST elevation on ECG) • Effusive constrictive pericarditis: Mixed clinical picture. Main clue is elevated JVP clinically & right atrial pressure on ECHO in spite of removal of pericardial fluid
<p>Essential tests:</p> <ul style="list-style-type: none"> • Chest X-ray • ECG • Echocardiogram 	<p style="text-align: center; background-color: #00bcd4; color: white; padding: 2px 5px; font-weight: bold;">INVESTIGATION</p> <p>Desirable:</p> <ul style="list-style-type: none"> • Cardiac enzymes • CT/MRI of Thorax • Pericardiocentesis • Pericardial biopsy

DIAGNOSIS

SUSPICION OF PERICARDIAL TUBERCULOSIS

MANAGEMENT

<p style="text-align: center; font-weight: bold; font-size: 10px;">TREATMENT</p> <ul style="list-style-type: none"> • Antitubercular therapy is advised as per NTEP • Steroids are recommended in large pericardial effusions, prominent pleocytosis & pericardial fluid with high inflammatory markers or early constriction • Give Prednisolone 60 mg/day for 4 weeks, 30 mg/day for 4 weeks, 15 mg/day for 2 weeks & 5 mg/day for 1 week • Total duration of systemic steroids is 11 weeks 	<p style="text-align: center; font-weight: bold; font-size: 10px;">NON RESPONSE TO STEROIDS & ATT</p> <ul style="list-style-type: none"> • Should prompt a referral to a specialist center for confirmation of diagnosis • Should prompt an evaluation for alternative causes of effusio-constrictive pericarditis: viral infections, systemic lupus erythematosus, primary effusion lymphomas or pericardial malignancies • Non response of cardiac symptoms to anti-tuberculous therapy cardiac surgical evaluation may be required
---	--

ABBREVIATION

ADA: Adenosine Deaminase	CXR: Chest X-ray	JVP: Jugular Venous Pressure
ATT: Antituberculous Therapy	ECG: Electrocardiogram	NTEP: National Tuberculosis Elimination Programme
	ECHO: Echocardiogram	TB: Tuberculosis

REFERENCES

1. National TB Elimination Programme, Central TB Division. Training Modules for Programme Managers & Medical Officers. Ministry of Health & Family Welfare, Government of India accessed at <https://tbcindia.gov.in/index1.php?lang=1&level=1&sublinkid=5465&lid=3540> Last access on 15 March, 2022.
2. Guidelines for programmatic management of drug resistant tuberculosis in India March 2021. National TB Elimination Programme, Central TB Division, Ministry of Health & Family Welfare, Government of India <https://tbcindia.gov.in/showfile.php?lid=3590> Last access on 15 March, 2022.
3. Nahid P, Dorman SE, Alipanah N, et al. Official American Thoracic Society/Centers for Disease Control and Prevention/Infectious Diseases Society of America Clinical Practice Guidelines: Treatment of Drug-Susceptible Tuberculosis. Clin Infect Dis. 2016 Oct 1;63(7):e147-e195. doi: 10.1093/cid/ciw376. Epub 2016 Aug 10.

This STW has been prepared by national experts of India with feasibility considerations for various levels of healthcare system in the country. These broad guidelines are advisory, and are based on expert opinions and available scientific evidence. There may be variations in the management of an individual patient based on his/her specific condition, as decided by the treating physician. There will be no indemnity for direct or indirect consequences. Kindly visit our web portal tbc.icmr.org.in for more information.

© Indian Council of Medical Research and Department of Health Research, Ministry of Health & Family Welfare, Government of India.

© 2024 JEFI

S222