

STANDARD TREATMENT WORKFLOW (STW)

Bacterial Skin Infections

Binod K Khaitan, Deepika Pandhi, Ananta Khurana, Dipankar De, Rahul Mahajan, Renu George, Vishal Gupta

¹All India Institute of Medical Sciences, New Delhi; ²University College of Medical Sciences, New Delhi; ³Dr. Ram Manohar Lohia Hospital, New Delhi; ⁴Postgraduate Institute of Medical Education and Research, Chandigarh; ⁵Postgraduate Institute of Medical Education and Research, Chandigarh; ⁶ Christian Medical College, Vellore; ⁷All India Institute of Medical Sciences, New Delhi

CORRESPONDING AUTHOR

Dr Binod K Khaitan, Department of Dermatology, All India Institute of Medical Sciences, New Delhi
Email: binodkhaitan@hotmail.com

CITATION

Khaitan BK, Pandhi D, Khurana A, De D, Mahajan R, George R, Gupta V. Bacterial Skin Infections. Journal of the Epidemiology Foundation of India. 2024;2(1Suppl):S81-S82.

DOI: <https://doi.org/10.56450/JEFI.2024.v2i1Suppl.041>

This work is licensed under a Creative Commons Attribution 4.0 International License.

@The Author(s). 2024 Open Access

DISCLAIMER

This article/STW, was originally published by Indian Council of Medical Research (ICMR) under Standard Treatment Workflow. The reprinting of this article in Journal of the Epidemiology Foundation of India (JEFI) is done with the permission of ICMR. The content of this article is presented as it was published, with no modifications or alterations. The views and opinions expressed in the article are those of the authors and do not necessarily reflect the official policy or position of JEFI or its editorial board. This initiative of JEFI to reprint STW is to disseminate these workflows among Health Care Professionals for wider adoption and guiding path for Patient Care.

July/2022



Department of Health Research
Ministry of Health and Family Welfare, Government of India



icmr
INDIAN COUNCIL OF
MEDICAL RESEARCH
Serving the nation since 1951

Standard Treatment Workflow (STW)

BACTERIAL SKIN INFECTIONS

ICD-10-L01, L73.9, L08, L02, L03, A46, L00

GENERAL PRINCIPLES OF MANAGEMENT

Skin hygiene, advise on handwashing/ local hygiene, avoidance of oil application, adequate nutrition

For recurrent/ severe lesions: evaluate for nasal carriage, diabetes, underlying skin conditions (scabies, atopic dermatitis)

In immunocompromised/ diabetics: consider the need for gram negative coverage

1. IMPETIGO

CLINICAL FEATURES

Wet yellow brown crusts overlying red inflamed skin

- **Types** Non bullous (NBI; commoner), bullous (BI)
- **Affected age group** usually children
- **Common sites** Face (perinasal, perioral) > extremities; extensive with scabies/ atopic eczema

MANAGEMENT

- Topical antibiotics for 5 days
- Oral antibiotics for extensive involvement or numerous lesions, lymphadenopathy or in outbreaks to prevent transmission

2. ECTHYMA

CLINICAL FEATURES

- Black thick crust (eschar) with underlying ulcer & surrounding redness & edema

MANAGEMENT

- Treat with oral antibiotics for 7 days
- Gentle crust removal may be attempted after soaks with sterile saline; topical antibiotics over the exposed ulcer

3. FOLLICULITIS

CLINICAL FEATURES

Hair follicle centred pustule/ papule
Rule out non bacterial causes: oils, chemicals, waxing, epilation, occlusive dressing
RECURRENT FOLLICULITIS Recurrent infection or outbreak in multiple members of family may indicate nasal *Staphylococcus aureus* carriage or human-pet transmission

MANAGEMENT

- Topical antibiotics for 5 days
- Oral antibiotics for multiple lesions
- Anti-inflammatory: Paracetamol 500mg/ Ibuprofen 400mg SOS for pain relief

4. FURUNCLE

5. CARBUNCLE

6. CUTANEOUS ABSCESS

Department of Health Research
Ministry of Health and Family Welfare, Government of India

Standard Treatment Workflow (STW)
BACTERIAL SKIN INFECTIONS
ICD-10-L01, L73.9, L08, L02, L03, A46, L00

GENERAL PRINCIPLES OF MANAGEMENT

- Skin hygiene, advise on handwashing/ local hygiene, avoidance of oil application, adequate nutrition
- For recurrent/ severe lesions: evaluate for nasal carriage, diabetes, underlying skin conditions (scabies, atopic dermatitis)
- In immunocompromised/ diabetics: consider the need for gram negative coverage

1. IMPETIGO

CLINICAL FEATURES
Wet yellow brown crusts overlying red inflamed skin

- **Types** Non bullous (NBI; commoner), bullous (BI)
- **Affected age group** usually children
- **Common sites** Face (perinasal, perioral) > extremities; extensive with scabies/ atopic eczema

MANAGEMENT

- Topical antibiotics for 5 days
- Oral antibiotics for extensive involvement or numerous lesions, lymphadenopathy or in outbreaks to prevent transmission

2. ECTHYMA

CLINICAL FEATURES
• Black thick crust (eschar) with underlying ulcer & surrounding redness & edema

MANAGEMENT

- Treat with oral antibiotics for 7 days
- Gentle crust removal may be attempted after soaking with sterile saline; topical antibiotics over the exposed ulcer

3. FOLLICULITIS

CLINICAL FEATURES
Hair follicle centred pustule/ papule
Rule out non bacterial causes: oils, chemicals, waxing, epilation, occlusive dressing

RECURRENT FOLLICULITIS Recurrent infection or outbreak in multiple members of family may indicate nasal *Staphylococcus aureus* carriage or human-pet transmission

MANAGEMENT

- Topical antibiotics for 5 days
- Oral antibiotics for multiple lesions
- Anti-inflammatory: Paracetamol 500mg/ Ibuprofen 400mg SOS for pain relief

4. FURUNCLE

CLINICAL FEATURES Painful follicle centric nodule/ pus point/ impending bulla/ ulcer with marked surrounding erythema, edema and induration

5. CARBUNCLE

CLINICAL FEATURES Confluence of multiple closely spaced furuncles + pus draining from multiple follicular orifices
Commonly nape of neck > breasts, buttocks in uncontrolled diabetes

6. CUTANEOUS ABSCESS

CLINICAL FEATURES Painful, warm, red fluctuant skin swelling

MANAGEMENT

- SMALL**
 - Oral antibiotics +
 - Topical antibiotics to reduce contamination of surrounding skin
- LARGE**
 - **INCISION AND DRAINAGE**
Incision and drainage/ debridement
 - Ancillary antibiotics if systemic inflammatory signs, associated septic phlebitis, multiple/ large abscesses, prominent cellulitis & immunocompromised state
- HOSPITALIZATION AND IV TREATMENT FOR SEVERELY ILL PATIENTS**
 - Inj Ceftriaxone 2g BD OR Inj Amoxicillin-clavulanate 1.2gm TDS
 - Alternatively - Inj Clindamycin 600-900mg TDS



7. CELLULITIS

CLINICAL FEATURES Acute spreading infection of skin involving subcutaneous tissue; Painful, red, tender, diffuse swelling mostly involving the limbs

8. ERYSIPELAS

CLINICAL FEATURES A more superficial, bright red, edematous, painful area with a clear demarcated edge; common sites: lower extremities > face. Often associated with lymphangitis and lymphadenopathy; broken skin/ portal of entry may be visualised

MANAGEMENT

CATEGORIZE DISEASE SEVERITY

- MILD**
 - Typical cellulitis/ erysipelas with no focus of purulence
 - Outpatient treatment with oral antibiotics
 - Elevation of affected area (to allow for dependent drainage); treatment of predisposing factors
 - Anti-inflammatory (Ibuprofen 400mg BD, Indomethacin 75mg BD)
- MODERATE**
 - Typical cellulitis/ erysipelas with systemic signs of infection
 - **MANAGEMENT**
 - **Hospitalization and parenteral antibiotics:**
 - Inj Ceftriaxone 2g BD OR Inj Amoxicillin-clavulanate 1.2gm TDS
 - Alternatively (allergic to penicillins) Inj Clindamycin 600-900mg IV TDS
- SEVERE**
 - With poor response to oral antibiotics, immunocompromised, signs of deeper infection like bullae, skin sloughing or systemic signs of infection like hypotension, or with organ dysfunction
 - **MANAGEMENT**
 - **Empiric broad spectrum IV antibiotic coverage**
 - Vancomycin + Piperacillin/ tazobactam
 - Surgical debridement
 - Sensitivity profile based modification of antibiotics

INVESTIGATIONS

1. Swabs for gram staining and pus culture are desirable
2. Blood cultures and biopsies are not routinely recommended, but useful with co-morbid conditions (malignancy on chemotherapy, immunocompromised states, animal bites etc.)

COMPLICATIONS
Subcutaneous abscesses, blistering (often haemorrhagic), ulceration, tissue necrosis, myositis, septicemia

9. STAPHYLOCOCCAL SCALDED SKIN SYNDROME

- Superficial peeling of skin due to toxin producing strains of staphylococcus
- Starts as tender and warm erythema and progresses to localised or generalised exfoliation with fever, malaise +/- dehydration and electrolyte disturbances
- Follows a local staphylococcal infection of either skin, throat, nose, umbilicus, or gut
- Bacteria cannot be demonstrated from blisters (cultures from original site may be positive)
- Treatment: preferably in-patient
- Mild cases: oral anti-staphylococcal antibiotics; severe cases: IV antibiotic
- Consider methicillin resistant *Staphylococcus aureus* (MRSA) coverage
- Usually remits within a week in children, high mortality in adults

RED FLAGS

- Temperature >100.4 F, WBC >12,000 or < 4000/μL, heart rate > 90 bpm, or respiratory rate > 24/min may indicate sepsis
- Severe pain followed by deceptive absence may indicate necrotising fasciitis
- Dark discoloration of overlying skin

PHARMACOTHERAPY

ANTIBIOTICS FOR SKIN AND SOFT TISSUE INFECTIONS

- PREFER β-LACTAMS**
 - Amoxicillin 500mg TDS (25-50 mg/kg/day)
 - Cloxacillin 500mg QID (50mg/kg/day)
 - Cephalixin 250-500mg QID (25-50 mg/kg/day)
 - Amoxicillin clavulanate combination: 625mg TDS
- IF ALLERGIC TO PENICILLINS**
 - Erythromycin 500mg QID (40 mg/kg/day)
 - Clindamycin: 300-600mg BD/TID (20mg/kg/day)

FOR NASAL CARRIERS
2% Mupirocin ointment for 5 days a month

TOPICAL ANTIBIOTICS

- Mupirocin cream 2%
- Fusidic acid cream 2%
- Framycetin cream 1%

IN ALL PATIENTS SUSPECT THE NEED FOR MRSA COVERAGE IF:

- Poor immune status
- Severe systemic signs
- MRSA infection elsewhere
- If no improvement in 48-72 hours
- Penetrating trauma

ORAL ANTIBIOTICS FOR SUSPECTED OR CONFIRMED MRSA INFECTION

- Cotrimoxazole 2 DS tablets BD
- Doxycycline 100 mg BD
- Minocycline 200 mg BD
- Linezolid 600 mg BD

IV ANTIBIOTICS FOR MRSA

- Vancomycin: 15 mg/kg BD
- Linezolid: 600 mg BD
- Clindamycin: 600-900 mg TDS

ANTIBIOTIC SUSCEPTIBILITY PATTERNS MAY VARY WITH REGION AND TIME

This STW has been prepared by national experts of India with feasibility considerations for various levels of healthcare system in the country. These broad guidelines are advisory, and are based on expert opinions and available scientific evidence. There may be variations in the management of an individual patient based on his/her specific condition, as decided by the treating physician. There will be no indemnity for direct or indirect consequences. Kindly visit the website of DHR for more information: (stw.icmr.org.in) for more information.

©Department of Health Research, Ministry of Health & Family Welfare, Government of India.